

# ESMO Meeting 2020 Late Breaking Abstract

## Title:

## 3-Years follow-up results of double-blind randomized phase II comparing concurrent high-dose cisplatin chemo-radiation plus Debio 1143 (xevinapant) or placebo in high-risk patients with locally advanced squamous cell carcinoma of the head and neck

## Authors:

J. Bourhis<sup>1</sup>, XS. Sun<sup>2</sup>, C. Le Tourneau<sup>5</sup>, Y. Pointreau<sup>3</sup>, C. Sire<sup>4</sup>, MC. Kaminsky-Forret<sup>7</sup>, A. Coutte<sup>6</sup>, M. Alfonsi<sup>8</sup>, P. Boisselier<sup>9</sup>, L. Martin<sup>10</sup>, JP. Delord<sup>11</sup>, F. Clatot<sup>12</sup>, J. Biau<sup>13</sup>, F. Rolland<sup>14</sup>, K. Gollmer<sup>15</sup>, S. Szyldergemajn<sup>15</sup>, C. Even<sup>16</sup> and Y. Tao<sup>16</sup>.

<sup>1</sup>CHUV, Lausanne, Switzerland

<sup>2</sup>Hôpital Nord Franche-Comté, Montbéliard, France

<sup>3</sup>Centre Jean Bernard, Le Mans, France

<sup>4</sup>Centre Hospitalier de Bretagne Sud - Hôpital du Scorff, Lorient, France

<sup>5</sup>Institut Curie, Paris, France

<sup>6</sup>C.H.U. Sud Amiens, Amiens, France

<sup>7</sup>L'Institut de Cancérologie de Lorraine (ICL) Alexis Vautrin, Vandoeuvre-lès-Nancy, France

<sup>8</sup>Institut Sainte Catherine, Avignon, France

<sup>9</sup>ICM - Val d'Aurelle, Montpellier, France

<sup>10</sup>Centre Guillaume le Conquérant, Le Havre, France

<sup>11</sup>Institut Claudius Regaud - Oncopole, Toulouse, France

<sup>12</sup>Centre Henri Becquerel, Rouen, France

<sup>13</sup>Centre Jean Perrin, Clermont-Ferrand, France

<sup>14</sup>Institut de Cancérologie de l'Ouest (ICO) René Gauducheau, Saint-Herblain, France

<sup>15</sup>Debiopharm International SA, Lausanne, Switzerland

<sup>16</sup>Institute Gustave Roussy, Villejuif, France

## Background:

This was a double-blind, randomized phase 2 study of 96 patients with poor prognosis locally advanced, squamous cell carcinoma of the head and neck (LA-SCCHN). Previously reported data showed significant improvements of Debio1143 versus placebo in addition to chemoradiation (CRT) for the primary endpoint of locoregional control (LRC) rate at 18 months, and demonstrated that addition of Debio 1143 was feasible, safe and did not compromise backbone therapy. Here we report up-dated data on PFS and overall survival (OS).

## Methods:

Patients, stratified by node involvement, tumor localization and HPV-16 status, were randomized (1:1) to receive Debio 1143 or matching placebo at 200 mg/day, oral once daily D1-14 q3w (3 cycles) when added to standard high-dose cisplatin CRT. PFS and OS were analyzed using a Cox model in the intention-to-treat population. Median and survival rates at 36 months were estimated by the Kaplan-Meier method.

## Results:

As of 21 Jul 2020, the median follow-up was 33 months. Debio 1143 combined with CRT showed a statistically significant improvement in OS vs placebo by reducing the risk of mortality by 51% (HR=0.49, [95%CI: 0.26-0.92], P=0.0261). The 3 years OS rate was 66% (95% CI: 49-78) in the Debio 1143 arm versus 51% (95% CI: 34-65) in the placebo arm; the median OS is not reached with Debio 1143 vs. 36.1 months with placebo (95%CI: 21.8-46.7). Statistically significant improvement in PFS was demonstrated, reducing the risk of disease progression or death by 66% (HR=0.34 [95%CI, 0.17-0.68], p=0.0023) and improving probability of PFS at 36 months to 72% in the Debio 1143 arm compared to 36% in the placebo arm.

The predictable and manageable safety profile observed with Debio 1143 + CRT after 2 years remained unchanged after 3 years.

## Conclusions:

These results with extended follow-up confirm those previously reported, showing now in addition a statistically and clinically significant OS benefit by adding Debio 1143 to standard CRT. The confirmatory phase 3 Trilynx study (EudraCT Number: 2020-000377-25) is ongoing.

**EudraCT Number: 2013-000044-25; ClinicalTrials.gov Identifier: NCT02022098**

